

# Misrepresentation in District Voting

Yoram Bachrach  
Microsoft Research  
Cambridge, United Kingdom  
yobach@microsoft.com

Yoad Lewenberg  
The Hebrew University of  
Jerusalem, Israel  
yoadlew@cs.huji.ac.il

Omer Lev  
University of Toronto  
Toronto, Canada  
omerl@cs.toronto.edu

Yair Zick  
Carnegie Mellon University,  
USA  
yairzick@cs.cmu.edu

## ABSTRACT

Voting systems in which voters are partitioned to districts encourage accountability by providing voters an easily identifiable district representative, but can result in a selection of representatives not representative of the electorate’s preferences. In some cases, a party may have a majority of the popular vote, but lose the elections due to districting effects.

We define the *Misrepresentation Ratio* which quantifies the deviation from proportional representation in a district-based election, and provide bounds for this ratio under various voting rules. We also examine probabilistic models for election outcomes, and provide an algorithm for approximating the expected Misrepresentation Ratio under a given probabilistic election model. Finally, we provide simulation results for several such probabilistic election models, showing the effects of the number of voters and candidates on the misrepresentation ratio.

## 1. INTRODUCTION

A voting system is a method by which voters choose between several alternatives and their opinions are aggregated, ultimately choosing a winner (or winners). Democratic countries, in principle, aim to have a representative outcome, by having a legislature roughly representative of the public’s beliefs, and in some countries, by having the chief executive elected directly by the public.

However, many democracies use a *district based* system for the selection of their legislature (most prominently, the Westminster system and the US system). In district based schemes, voters are divided into geographically based districts<sup>1</sup>, and each one selects a representative to the legislature. The selection mechanisms differ: Westminster and US systems use plurality, France uses plurality with runoff, while Australia uses STV. Typically, candidates in each constituency are members of political parties, and some systems have the majority party form the executive (others, such as

<sup>1</sup>While we use the term “district”, other terms include “electoral district” (US), “riding” (Canada), and “constituency” (UK).

**Appears at:** 3rd Workshop on Exploring Beyond the Worst Case in Computational Social Choice. Held as part of the 15th International Conference on Autonomous Agents and Multiagent Systems. May 9th-10th, 2016. Singapore.

the US, have a similar process for selecting the chief executive).

Single-member district elections provide voters a single and easily identifiable district representative, encouraging service and accountability. However, the proportion of seats in the legislature belonging to a party may be very different from the proportion of voters supporting that party in the overall population; this is known as the “referendum paradox” [33]. The disparity between the popular vote and the district vote has been a source of contention in US elections; by redistricting constituencies (“gerrymandering”), political parties have manipulated the elections [25, 10]; indeed, the US Voter Rights Act of 1965 includes several provisions that require change in congressional districts in several states to be approved by federal authorities [36].

Moreover, such a discrepancy is caused not only by gerrymandering, but is built into district-based mechanisms. In the US presidential election of 1876, the losing candidate, Samuel Tilden, got 6% more votes than the winner, Rutherford Hayes, an occurrence that happened twice since. The electoral college, through which the president is elected, displays the problem even more acutely; in the US 1992 presidential election a candidate that garnered 18.9% of the vote, Ross Perot, was not represented at all. In the UK system, in 1951 the Conservative party lost the popular vote to the Labour party while still winning a strict parliamentary majority; in general, while no party has received a majority of the popular vote in a British election since 1931, all but two elections resulted in a parliamentary majority for one of the parties. Similar scenarios have unfolded in Canada, Australia and elsewhere.

Such problems may also arise in any multi-level decision making process. So, if an organization or a sensor analysis system (e.g., an automated car), attempts to decide on its next step based on inputs from sub-systems employing their own decision making processes (e.g., each sensor family is a district, and sensors are “voting” between themselves), they may also encounter such a problem, as a small amount of signals may cause the system to reach a wrong outcome.

Consider the following example. Two political parties, *A* and *B* compete in seven districts of equal size,  $D_1, D_2, \dots, D_7$ . Both parties run a candidate in every district, and the plurality voting rule is used to determine the winner. Now, suppose that in districts  $D_1, \dots, D_4$ , 60% of the vote goes to the representative of party *A*; on the other hand, in districts  $D_5, D_6$  and  $D_7$  the representatives of party *B* take 100% of the vote. The final vote tally shows party *A* wins

the election, even though party  $B$  has nearly twice as many voters!

As the example above shows, a minority (those in favor of party  $A$ ) may rule over the majority (those in favor of party  $B$ ). We refer to the outcome in this case as a *misrepresentation of the voters’ preferences*. District based electoral systems tend to be more “stable”: they tend to result in a smaller number of candidates, and thus less fragmentation of the parliamentary body. However, misrepresentation is an inherent byproduct of electoral stability; indeed, stability comes at the heavy price of potentially overriding the preferences of most voters.

One way to quantify the degree to which a system skews the true desires of the voters — as captured by the total support for each party — is to examine the ratio between the number of those who voted for the most popular party in general, and the number of voters who voted for the winner. When more people voted for the losing party than for the winning party this ratio is larger than one. The higher this ratio is, the more pronounced the misrepresentation effect.

#### *Our Contribution.*

We examine the issue of misrepresentation due to district-based systems under several prominent voting rules. We first provide a metric for quantifying the misrepresentation effect, both for elections with two parties and for elections with more than two parties, which we call the Misrepresentation Ratio. We provide bounds on the Misrepresentation Ratio for several voting rules, depending on the numbers of voters, candidates and districts. Finally, we provide some simulation results regarding the misrepresentation ratio in certain scenarios, to examine its values under various settings.

## 2. RELATED WORK

Voting district representability has been the topic of much public debate and research for two centuries, since the advent of redistricting in United States (“gerrymandering”) <sup>2</sup>

and in United Kingdom, redistricting due to an attempt to follow population changes in successive reform bills. Most of the research on these issues focused on historical [8, 7], sociological [29] and political science [10] issues. Particularly since the Voting Rights Act of 1965, much legal research has also dealt with district based system, though usually focusing on particular countries (mainly – though not solely – the US) [36, 25].

Political science research on such misrepresentation has mostly focused on trying to assess when it might occur, studying the history of elections or using statistic assumptions [38] and analysis on election data [16, 31, 32, 27, 28, 22]. This analysis, however, does not try to assess how bad the misrepresentation can get, and mostly focuses on plurality and its variants. A somewhat similar line of research tries to assess voter’s misrepresentation as distance of parliamentary seat allocation compared to pure representational settings [19, 17, 20, 15], applying concepts such as the Banzhaf

<sup>2</sup>Our analysis focused on how bad misrepresentation might be, rather than the issue of how restructuring districts may be used as a purposeful manipulation, designed to give an unfair advantage to a specific candidate, more in line with other forms of manipulation and control in elections [11, 13, 1, 3].

index and voting power. However, we deal with the different issue of quantifying the misrepresentation degree by examining the degree of support of the losing candidate.

The computational social choice community focused on manipulation by voters [40, 41, 35], and less on more institutional motivated manipulation. Some work has been done on control problems, in which a central authority may significantly influence results (e.g. [23] and the survey by [12]), including preliminary work on dividing voters into groups [9]. Optimal gerrymandering strategies have been studied in [34, 18]; however, these have mostly focused on 2 party scenarios, as in the US.

In establishing our bounds, we encounter the problem of finding the minimal score a candidate can win with, given a scoring rule; [39] studies a similar problem, but their results do not directly apply to our setting. Our problem is a special case of the bin packing problem with a constant number of types; this problem has been widely believed to be NP-hard, but recent advances [37, 26] have eventually established that it is in P [21].

## 3. PRELIMINARIES

We have a set of *voters*,  $V$ , and each voter  $i \in V$  has a preference order (without ties) over the *candidate set*  $C$ , denoted  $\prec_i$ . For every  $c, c' \in C$ , we say that  $i$  *prefers*  $c$  to  $c'$  if  $c' \prec_i c$ . We denote the set of linear preferences over  $C$  as  $\mathcal{L}(C)$ .

A *voting rule* is a function  $f : \mathcal{L}(C)^n \rightarrow C$ , whose input is a finite list of linear preferences over  $C$  (a *profile*) of size  $n$  (most voting rules are well defined for any  $n > 0$ ) and whose output is a candidate  $c \in C$ . Voting rules are often assumed to output a set of candidates, but since in our setting we are only interested in the winners of the election, the output of  $f$  is a single candidate ( $f$  may incorporate some fixed tie-breaking rule). Since the set of voters and their preferences is constant throughout the paper, we overload notation and define  $f$  over subsets of  $V$ : given a set  $Q \subseteq V$ , we let  $f(Q)$  be the output of  $f$  over the preferences of the voters in  $Q$ .

A voting rule is *neutral* if the outcome is invariant under candidate name changes. More formally, a voting rule  $f$  is neutral if for any two candidates  $c$  and  $c'$ , if every voter  $i \in V$  now ranks  $c$  in the position of  $c'$  and vice versa, the outcome of  $f$  is unchanged if the winner was neither  $c$  or  $c'$ , is  $c'$  if the winner was  $c$ , and is  $c$  if the winner was  $c'$ .

We say a voting rule  $f$  is *score-monotone* if it induces a score for every candidate  $c \in C$  (i.e., each candidate ends up with some real number, and the one with the maximal one is chosen), and the following holds: given two preference profiles  $R_1 = (\prec_i)_{i \in V}$  and  $R_2 = (\prec'_i)_{i \in V}$ , if for all  $a, b \in C \setminus \{c\}$  we have  $a \prec_i b \iff a \prec'_i b$ , and under  $R_1$ ,  $c$  is in a position that is no lower than its position in  $R_2$ , then  $c$ ’s score under  $R_1$  is at least as high as its score in  $R_2$ . These properties hold for the common voting rules which will be used in this paper.

**Scoring Rules:** A scoring rule is defined by a vector of scores,  $(\alpha_1, \dots, \alpha_m)$ , where  $\alpha_1 \geq \dots \geq \alpha_m = 0$ ,  $\alpha_1 > 0^3$  and  $m$  is the number of candidates. Given a voter  $i \in V$ , let  $rank_i(c)$  be the rank of candidate  $c$  in the preference order of  $i$ . Given a set of voters  $Q \subseteq V$ ,

<sup>3</sup>Assuming  $\alpha_m = 0$  is no loss of generality: if  $\alpha_m > 0$  score vector can be normalized by setting  $\alpha_j = \alpha_j - \alpha_m$ .

let  $score_c^\alpha(Q) = \sum_{i \in Q} \alpha_{rank_i(c)}$ . Given a set of voters  $Q \subseteq V$ , the output of  $Scoring_\alpha(Q)$  is a member of  $\arg \max_{c \in C} score_c^\alpha(Q)$  (if there are ties, we break them according to some tie-breaking rule). Many scoring rules are widely used. For example, the *plurality* rule, where  $\alpha_1 = 1$  and  $\alpha_j = 0$  for all  $j \geq 2$ , the *veto* rule, where  $\alpha_j = 1$  for all  $j < m$ , and  $\alpha_m = 0$ , both of which are specific examples of the family of  $k$ -approval scoring rules, in which  $\alpha_1 = \dots = \alpha_k = 1$  and  $\alpha_{k+1} = \dots = \alpha_m = 0$ . Another common scoring rule is the *Borda* rule, where  $\alpha_j = m - j$  for all  $j \in \{1, \dots, m\}$ .

**Copeland Rule:** Given a set of voters  $Q \subseteq V$ , we say that a candidate  $c$  beats  $c'$  in a pairwise election under  $Q$  if a majority of the members of  $Q$  prefer  $c$  to  $c'$ . For each candidate  $c \in C$ , we let  $score_c^{Cp}(Q)$  to be the number of candidates that  $c$  beats in a pairwise election, minus the number of candidates that beat  $c$ . The Copeland voting rule outputs  $Copeland(Q) = \arg \max_{c \in C} score_c^{Cp}(Q)$ .

We note that both scoring rules and Copeland have a natural notion of a candidate score, and were conceptualized as such in [6]. This will be instrumental in defining voting misrepresentation, as we do in the following section.

*Definition 1.* Given a voting rule  $f$  which induces a candidate score, we let  $score_f(c, Q)$  to be the score of  $c \in C$  under  $f$ , when the voter set is  $Q \subseteq V$ .

## 4. THE MISREPRESENTATION RATIO

We are interested in settings where the set of voters is partitioned into districts: these are  $z$  disjoint sets  $D_1, \dots, D_z$  whose union is  $V$ . The election winner under this model is determined by applying the voting rule  $f$  to  $D_1, \dots, D_z$ ; the candidate who wins the greatest number of districts is the winner (ties are broken using some tie-breaking rule).

*Definition 2.* Given a partition of voters into  $z$  districts  $\mathcal{D} = \{D_1, \dots, D_z\}$ , let  $w$  be the winner of the election when voters are partitioned into districts as per  $\mathcal{D}$ . The *misrepresentation ratio* is the ratio between the maximum score of any candidate under  $f$ , and the score of  $w$ . Formally:

$$MR(V, \mathcal{D}, f) = \frac{\max_{c \in C} score_f(c, V)}{score_f(w, V)}.$$

Note that  $MR(V, \mathcal{D}, f) \geq 1$ ; if  $MR(V, \mathcal{D}, f) = 1$  then the winner of the district elections completely captures the popular vote (as measured by  $f$ ). The higher  $MR(V, \mathcal{D}, f)$ , the less popular the winning candidate is in the eyes of the people; thus, district elections with a high misrepresentation ratio are ones where voters' preferences are not appropriately aggregated, due to the effects of district elections.

*Remark 1.* In this work we assume all districts have the same number of voters. Without any assumptions on the number of voters in each district, the worst case misrepresentation ratio (Definition 2) can be arbitrarily high: consider  $2\ell + 1$  districts. Suppose  $\ell + 1$  of the districts have only three voters, and  $\ell$  of them have  $M$  voters, where  $M$  is a very large number. There are two candidates,  $A$  and  $B$ ; candidate  $A$  wins all votes in the  $\ell$  districts where there are  $M$  voters, and candidate  $B$  wins 2 of the 3 votes in the

$\ell + 1$  districts holding three voters. Thus, the total number of votes for  $B$  is  $\mathcal{O}(\ell)$ , and the total number of votes for  $A$  is  $\mathcal{O}(M\ell)$ , resulting in an arbitrarily high misrepresentation. Our results can be extended by incorporating an additional parameter:  $\max_{D, D' \in \mathcal{D}} \frac{|D|}{|D'|}$ ; however, in the interest of space and clarity, we assume that districts are of equal size.

## 5. BOUNDS ON THE MISREPRESENTATION RATIO

In what follows, we establish upper bounds on the *worst-case misrepresentation ratio*; we show that our bounds are tight, i.e. that there exist district elections where the bound holds.

Furthermore, we always refer to the district voting instance  $\langle V, C, (\prec_i)_{i \in V}, \mathcal{D}, f \rangle$  as one that maximizes  $MR$ , where  $|C| = m$ ,  $|\mathcal{D}| = z$ , for all  $D \in \mathcal{D}$ :  $|D| = n$ , and  $f$  is some neutral score-monotone rule. We assume that  $w \in C$  is the winner of the district elections; that is,  $w$  won a plurality of the districts. We mark  $d(c)$  for  $c \in C$  as the number of districts won by candidate  $c$ . We assume that in the case of a tie for first place in a district, ties are broken against the district election winner; moreover, the district election winner must win a strict majority of the districts.

Intuitively, in order to establish our bound, we want to create the worst possible election. Such an election would have the candidate  $w$  win by as small a margin as possible, with some other candidate  $p \neq w$  being as popular as possible while losing a majority of the districts.

Given a score inducing voting rule  $f$ , we let  $L_n(f)$  be the maximal score that a candidate can get while still losing an  $n$  voter election, and  $M_n(f)$  be the maximal score that a candidate can get and win an  $n$  voter election. For example,  $L_n(\text{Plurality}) = \lceil \frac{n}{2} \rceil - 1$ , and  $M_n(\text{Plurality}) = n$ . The following lemma offers some insights regarding the number of districts won by each candidate in an instance maximizing  $MR$ .

LEMMA 1. *If  $f$  is a score-monotone, neutral rule then there is a district voting instance maximizing  $MR$  such that:*

1. *if  $w$  does not win the election in a district  $D$ , then  $w$  is ranked last by all voters in  $D$ .*
2. *If the candidate  $p$  wins in  $D$ , then it is ranked first by all voters in  $D$ ; if not and the winner is not  $w$ , then the score of  $p$  from  $D$  is  $L_n(f)$ .*
3.  *$d(p) = d(w) - 1$  if  $m > 2$  or  $z$  is odd, otherwise  $d(p) = d(w) - 2$ .*

PROOF. If  $w$  loses the election in a district  $D$ , let  $i \in D$  be a voter who did not rank  $w$  last; we can simply swap the candidate that was ranked last by  $i$  with  $w$  in  $i$ 's ranking. This will result in a voter profile where the score of  $w$  is weakly lower, and the score of every other candidate is weakly higher (by the score monotonicity and neutrality of  $f$ ). Furthermore, if  $p$  wins the election in  $D$ , but is not ranked first by some voter  $i \in D$ , swapping  $p$  with the candidate ranked first by  $i$  (this is not  $w$  by the previous observation) will result in  $p$  having a weakly higher score. Whenever  $p$  loses the election in a district, its score from that district is lower than the score that it could have obtained by winning a district, but as we wish to maximize it

(and we do not care for the score of candidates other than  $w$  and  $p$ ),  $p$  should score  $L_n(f)$ .

To show that  $d(c) \leq d(p)$  for all  $c \neq w$ , suppose there is a candidate  $c \neq w, p$  for which  $d(c) > d(p)$ . Changing districts who voted for  $c$  to vote for  $p$  will only improve  $p$ 's score while not changing  $w$  as winner; thus,  $d(c) \leq d(p)$  for all  $c \neq p, w$ .

For 2 candidates and an even number of districts, the difference between the number of districts must be at most 2; if it is more, we can take districts voting for  $w$  and change them to  $p$ , improving  $p$ 's score and hurting  $w$  (while having  $w$  remain the election winner). For 2 candidates, if the number of districts is odd we do a similar process, reaching a difference of 1.

Now, suppose  $w$  has won  $q$  more districts than  $p$ , and  $q > 2$ . In this case, we can take some district where  $w$  won and make  $p$  the winner in that district. That would strictly improve  $p$ 's score, reduce  $w$ 's score and maintain  $w$  as the overall election winner. This establishes that  $d(w) \leq d(p) + 2$ . Now, we wish to show that if  $m \geq 3$  then  $d(w) = d(p) + 1$ . If  $d(c) = 0$  for all  $c \neq w, p$  then we are back in the two candidate case which we have previously covered, and if  $z$  is even, we can give one of  $w$ 's district to  $c$ . If there is some district  $D$  won by some  $c \neq w, p$  and  $d(w) = d(p) + 2$  then we can make  $p$  win  $c$ 's district, weakly improving its score while maintaining  $w$ 's score at the same level, and keeping  $w$  as the district election winner.  $\square$

## 5.1 Scoring Rules

We begin our investigation by bounding  $MR$  when  $f$  is a scoring rule. We begin with a simple technical lemma.

LEMMA 2. *For every scoring rule  $f = \text{Scoring}_\alpha$ , where  $\alpha = (\alpha_1, \dots, \alpha_m)$ , if  $\alpha_1 > \alpha_2$ ,  $L_n(f) = \alpha_1(\lceil \frac{n}{2} \rceil - 1) + \alpha_2(\lfloor \frac{n}{2} \rfloor + 1)$ . If  $\alpha_1 = \alpha_2$ , denote  $\alpha'$  as the maximal  $\alpha_i$  such that  $\alpha_i < \alpha_1$ , then  $L_n(f) = (n - 1)\alpha_1 + \alpha'$ .*

PROOF. We shall consider some candidate  $c \in C$  which we make the winner, while giving candidate  $p \in C$  the maximal score for a runner up. First, showing that there indeed can be a winner with a higher score, if  $\alpha_1 > \alpha_2$ , if  $\lceil \frac{n}{2} \rceil - 1$  voters give  $\alpha_1$  to  $p$  and  $\alpha_2$  to  $c$ , while the rest of the voters give  $\alpha_1$  to  $c$  and  $\alpha_2$  to  $p$ , we reach the desired score for  $p$  while  $c$  is the winner. If  $\alpha_1 = \alpha_2$ , for  $n - 1$  voters giving  $\alpha_1$  or  $\alpha_2$  to both  $p$  and  $c$ , and the final voter giving  $\alpha_1$  points to  $c$  and  $\alpha'$  points to  $p$ , we again reach the desired score while  $c$  is the winner.

Now, we shall show no higher score is possible. For the case  $\alpha_1 = \alpha_2$  it is obvious: the only higher score possible is  $n\alpha_1$ , which is the maximal one, hence the winner's. For  $\alpha_1 > \alpha_2$ , suppose there is a better  $L_n(f)$ , which is described as  $z_1\alpha_1 + z_2\alpha_2 + \dots + z_m\alpha_m$ . If  $z_1 \leq \lceil \frac{n}{2} \rceil - 1$ , then this sum is smaller than that suggested in the lemma. If  $z_1 > \lceil \frac{n}{2} \rceil - 1$ , this means the winner  $c$  has a score of at most  $\alpha_1(\lceil \frac{n}{2} \rceil - 2) + \alpha_2(\lfloor \frac{n}{2} \rfloor + 1)$ . This score needs to be higher than  $z_1\alpha_1 + z_2\alpha_2 + \dots + z_m\alpha_m$ , but it is lower than the lemma's suggested  $L_n(f)$ , reaching a contradiction.  $\square$

COROLLARY 1. *If  $f = \text{Scoring}_\alpha$  then  $(z + 1)L_n(f) \geq M_n(f)$  for any  $z > 1$ .*

PROOF. Given a vector  $\alpha = (\alpha_1, \dots, \alpha_m)$  we have that  $M_n(f) = n\alpha_1$ . Now, by Lemma 2, we know that

$$L_n(\text{Scoring}_\alpha) \geq \left( \left\lceil \frac{n}{2} \right\rceil - 1 \right) \alpha_1,$$

from which the claim trivially follows.  $\square$

The following lemma tells us the number of districts that must be won by the winning candidate in a district election that maximizes  $MR$ .

LEMMA 3. *Let  $\langle V, C, (\prec_i)_{i \in V}, \mathcal{D}, f \rangle$  be a district voting instance which maximizes  $MR$ , where  $f$  is a scoring rule.  $z$  (number of districts) can be written as  $\ell m + r$  ( $\ell, r \in \mathbb{N} \cup \{0\}$  and  $r < m$ ). There is an  $MR$  maximizing instance where the number of districts that  $w$  wins is at most  $\ell + 2$ , and every other candidate wins at least  $\ell - 1$  districts.*

PROOF. Recall that  $d(c)$  be the number of districts that each candidate  $c \in C$  wins. In order to win the district election,  $w$  needs to win a plurality of the districts; that is,  $d(w) \geq d(c) + 1$  for all  $c \in C \setminus \{w\}$ . Now, let us denote the score of  $w$  from winning a district as  $x_w$ ; by Lemma 1  $w$  gets no votes from all districts where it loses (the case where there are two candidates and an even number of districts can be handled similarly, and is omitted due to space constraints). Now,  $p$  gets a score of  $x_p$  from a district that  $w$  wins, a score of  $M_n(f)$  from a district that it wins, and a score of  $L_n(f)$  from a district that some other candidate wins. Putting it all together we have:

$$\frac{x_p d(w) + M_n(f) d(p) + L_n(f) \left( \sum_{c \in C \setminus \{w, p\}} d(c) \right)}{x_w d(w)} = \frac{x_p}{x_w} + \frac{M_n(f) d(p) + L_n(f) \left( \sum_{c \in C \setminus \{w, p\}} d(c) \right)}{x_w d(w)}.$$

Since  $d(w) = d(p) + 1$ ,  $\sum_{c \in C \setminus \{w, p\}} d(c) = z - 2d(w) + 1$ , hence our equation is

$$\frac{x_p + M_n(f) - 2L_n(f)}{x_w} + \frac{(z + 1)L_n(f) - M_n(f)}{x_w d(w)}. \quad (1)$$

We are interested in maximizing (1) as a function of  $d(w)$ ; the derivative of (1) with respect to  $d(w)$  is

$$\frac{(z + 1)L_n(f) - M_n(f)}{x_w} \cdot \frac{1}{-d(w)^2}$$

which is positive if and only if  $(z + 1)L_n(f) < M_n(f)$ . According to Corollary 1,  $(z + 1)L_n(f) > M_n(f)$  then (1) is maximized when  $d(w)$  is as small as possible. If  $r = 0$  then  $d(w) = \ell + 1$  while every other candidate gets  $\ell$  (if  $d(w) \leq \ell$  more than one candidate will have such a score). If  $r \geq 1$ , then  $d(w)$  must be more than  $\ell + 1$ , as more than one candidate must get that score, so  $d(w) = \ell + 2$  and other candidates can get  $\ell + 1$  districts or less.  $\square$

The following lemma discusses the score that  $w$  receives in districts that it wins.

LEMMA 4. *For every scoring rule  $f$ , if  $n \gg z$ ,  $z > 3$  (i.e., number of voters in each district is far larger than the number of districts), the score given to  $w$  in districts in which it wins must be the minimal possible score needed to win a district in an instance maximizing  $MR$ .*

PROOF. Note that this is not trivial – it is not obvious that we do not wish to increase  $w$ 's score to allow  $p$  to receive a higher score as a 2nd place candidate.

For simplicity, we shall assume  $n$  is an odd number and  $\alpha_1 > \alpha_2$ . The proof is equivalent in the other cases. We can also assume  $m > 2$ , as otherwise this is already set by Lemma 1.

From Equation 1, we know that we seek to maximize

$$\frac{x_p + M_n(f) - 2L_n(f)}{x_w} + \frac{(z+1)L_n(f) - M_n(f)}{x_w d(w)}.$$

We know  $M_n(f) = n\alpha_1$  and  $L_n(f) = \alpha_1(\lceil \frac{n}{2} \rceil - 1) + \alpha_2(\lfloor \frac{n}{2} \rfloor + 1)$  from Lemma 2, hence we wish to maximize:

$$\begin{aligned} & \frac{x_p + n\alpha_1 - (n-1)\alpha_1 - (n+1)\alpha_2}{x_w} + \\ & \frac{(z+1)(\lceil \frac{n}{2} \rceil \alpha_1 + \lfloor \frac{n}{2} \rfloor \alpha_2) - n\alpha_1}{x_w d(w)} \\ = & \frac{x_p + \alpha_1 - (n+1)\alpha_2}{x_w} + \frac{(z+1)(\lceil \frac{n}{2} \rceil \alpha_1 + \lfloor \frac{n}{2} \rfloor \alpha_2) - n\alpha_1}{x_w d(w)}. \end{aligned}$$

Now, whatever  $x_w$  is, we wish to maximize  $x_p$  as long as  $x_w > x_p$ . For a particular scoring rule  $f$ ,  $x_p = x_w - s_n(f)$  for some function  $s_n^4$ . Therefore, looking at the derivative of our equation according to  $x_w$ , we get

$$\begin{aligned} & \frac{1}{x_w^2} \left( s_n(f) - \alpha_1 + (n+1)\alpha_2 - \right. \\ & \left. \frac{(z+1)(\lceil \frac{n}{2} \rceil \alpha_1 + \lfloor \frac{n}{2} \rfloor \alpha_2) + n\alpha_1}{d(w)} \right) \\ = & \frac{1}{x_w^2} \left( s_n(f) + \alpha_1 \frac{n-d(w)-(z+1)\lceil \frac{n}{2} \rceil}{d(w)} + \right. \\ & \left. \alpha_2 \frac{d(w)(n+1)-(z+1)\lfloor \frac{n}{2} \rfloor}{d(w)} \right). \end{aligned}$$

As  $n \gg z$  and  $z > 3$ , this is always negative as the  $\alpha_1$  and  $\alpha_2$  components overwhelms all others (since  $s_n(f) < n\alpha_1$ ), hence  $x_w$  needs to be minimal in size.  $\square$

LEMMA 5. *For  $k$ -approval voting rules, the minimal score a winner can get in an election is  $\lfloor \frac{nk}{m} \rfloor + 1$  if  $m \lfloor \frac{nk}{m} \rfloor + 1 = nk$ . Otherwise, it is  $\lceil \frac{nk}{m} \rceil + 1$ .*

PROOF. Since  $nk$  points are allocated by voters, a winner must have at least a score of  $\lfloor \frac{nk}{m} \rfloor + 1$ . If  $m \lfloor \frac{nk}{m} \rfloor + 1 = nk$ , then it is possible that  $w$  gets  $\lfloor \frac{nk}{m} \rfloor + 1$  votes, while all other candidates get  $\lfloor \frac{nk}{m} \rfloor$ . If  $m \lfloor \frac{nk}{m} \rfloor + 1 < nk$ , there is no allocation of  $nk$  points such that only a single candidate gets  $\lfloor \frac{nk}{m} \rfloor + 1$ , hence the winner will need to get one more point, with the rest of the points to be allocated among the other candidates, with none of them getting more than  $\lceil \frac{nk}{m} \rceil$  votes.  $\square$

Armed with Lemmata 1 – 5, we now proceed to analyze specific voting rules.

THEOREM 1. *Suppose that  $n = qm + s$  and  $z = \ell m + r$ , where  $q, s, \ell, r \in \mathbb{N} \cup \{0\}$  and  $s, r < m$ ; then  $MR(V, \mathcal{D}, \text{Plurality})$  is at least*

$$1 + \frac{n - 2 \lceil \frac{n}{2} \rceil + 1}{q + 2} + \frac{(z+1)(\lceil \frac{n}{2} \rceil - 1) - n}{(\ell + 2)(q + 2)}$$

and at most

$$1 + \frac{n - 2 \lfloor \frac{n}{2} \rfloor + 1}{q + 1} + \frac{(z+1)(\lfloor \frac{n}{2} \rfloor - 1) - n}{(\ell + 1)(q + 1)}.$$

In particular,  $MR(V, \mathcal{D}, \text{Plurality})$  is in  $\Theta(m^2)$ .

<sup>4</sup>Obviously,  $x_w - n \max_i(\alpha_i - \alpha_{i+1}) \leq x_p$ , as  $x_p$  can always be ranked below  $x_w$ . On the other hand,  $x_p \leq x_w - \min_i(\alpha_i - \alpha_{i-1})$ .

PROOF. By Lemma 3 and Corollary 1, we know that  $w$  wins either  $\ell + 1$  or  $\ell + 2$  districts. Plugging in the values  $L_n(\text{Plurality}) = \lceil \frac{n}{2} \rceil - 1$  and  $M_n(\text{Plurality}) = n$  into (1), we have that  $MR(V, \mathcal{D}, \text{Plurality})$  equals

$$\frac{x_p + n - 2 \lceil \frac{n}{2} \rceil + 2}{x_w} + \frac{(z+1)(\lceil \frac{n}{2} \rceil - 1) - n}{x_w d(w)}$$

( $x_w$  and  $x_p$  denote the score of  $w$  and  $p$  (respectively) when  $w$  wins a district). We are left just with determining the value of  $x_p$  and  $x_w$ . For reasons similar to the ones detailed in Lemma 1, it holds that  $x_p = x_w - 1$ .

Due to Lemma 4,  $MR(V, \mathcal{D}, \text{Plurality})$  is maximized when  $x_w$  is minimal; We have that  $w$  receives  $q + 2$  votes if  $s \geq 2$ , and  $q + 1$  otherwise. Plugging this into (1) we obtain the desired result.

The second expression in the upper bound of Theorem 1,  $\frac{n - 2 \lceil \frac{n}{2} \rceil + 1}{q + 1}$  can be upper bounded by  $\frac{1}{q + 1}$ , which is at most  $\frac{m}{n}$ . Thus, if the number of voters dominates the number of candidates, this expression has little effect on  $MR$ . The second expression can be upper bounded as follows

$$\frac{(z+1)(\lceil \frac{n}{2} \rceil - 1) - n}{(\ell + 1)(q + 1)} \leq \frac{zn}{(\ell + 1)(q + 1)} \leq m^2$$

A similar lower bound of  $\Omega(m^2)$  can be shown as well, which concludes the proof.  $\square$

Note that tightness is achieved as our constructed expressions were dependent on particular voting profiles (as described in Lemma 1), and hence carry on to these expressions.

As some parliamentary systems require not a plurality of districts to become a winner, but a majority, we also note the  $MR$  in these cases.

COROLLARY 2. *If the number of districts needed for a victory is above 50%,  $MR$  for plurality is  $\Theta(m)$ .*

PROOF. Suppose that  $n = qm + s$  for  $q, m \in \mathbb{N}$ ,  $s < m$ . As  $d(w)$  in our expression is now  $\lfloor \frac{n}{2} \rfloor + 1$ , and thanks to Lemma 4 we know we should minimize  $x_w$ . Hence,  $w$  receives  $q + 2$  votes if  $s \geq 2$ , and  $q + 1$  otherwise. Thanks to Lemma 1  $d(p) = \lceil \frac{n}{2} \rceil - 1$ , and its score in the other districts is  $x_w - 1$ . The formula turns out to be, for  $s \geq 2$  (very similar figure for  $s \leq 1$ ):

$$\frac{q + 1}{q + 2} + \frac{n(\lceil \frac{n}{2} \rceil - 1)}{(q + 2)(\lfloor \frac{n}{2} \rfloor + 1)} \xrightarrow{n \rightarrow \infty} m.$$

$\square$

THEOREM 2. *Suppose that the number of districts is expressed as  $z = \ell m + r$ , where  $\ell, r \in \mathbb{N} \cup \{0\}$  and  $r < m$ ; then  $MR(V, \mathcal{D}, k\text{-approval})$  for  $k > 1$  is at least*

$$1 + \frac{n(z + \ell + 2) - (z - \ell + 1)}{(\ell + 2)(\lfloor \frac{nk}{m} \rfloor + 1)}$$

and at most

$$1 + \frac{n(z + \ell + 1) - (z - \ell)}{(\ell + 1)(\lfloor \frac{nk}{m} \rfloor + 1)}.$$

PROOF. By Lemma 3 and Corollary 1, we know that  $w$  wins either  $\ell + 1$  or  $\ell + 2$  districts. Plugging in the values  $L_n(k\text{-approval}) = n - 1$  and  $M_n(k\text{-approval}) = n$  into (1), we have that  $MR(V, \mathcal{D}, k\text{-approval})$  equals  $\frac{x_p + 2 - n}{x_w} + \frac{kn - (z + 1)}{x_w d(w)}$

( $x_w$  and  $x_p$  denote the score of  $w$  and  $p$  (respectively) when  $w$  wins a district). As before, for reasons similar to the ones detailed in Lemma 1,  $x_p = x_w - 1$ .

Due to Lemma 4,  $MR(V, \mathcal{D}, k\text{-approval})$  is maximized when  $x_w$  is minimal; According to Lemma 5,  $x_w = \lfloor \frac{nk}{m} \rfloor + 1$  if  $m \lfloor \frac{nk}{m} \rfloor + 1 = nk$ . Otherwise, it is  $x_w = \lceil \frac{nk}{m} \rceil + 1$ . Plugging these values into the equation, we receive the desired result.  $\square$

One of the main challenges in computing a closed form formula for  $MR$  for general scoring rules is that one must first decide what is the minimal score that  $w$  can obtain while winning a district for a given score vector  $\alpha$ . This problem can be thought of as a *bin packing problem*: candidates can be thought of as bins, and the scores must be packed into them. It is only recently that a polynomial time algorithm has been proposed for bin packing problems with a constant number of types (also commonly referred to as the *one-dimensional cutting stock problem*) [21]. Thus, for general scoring rule we offer looser bounds on the number of votes needed to win:

Let  $S_\alpha = \sum_{j=1}^m \alpha_j$ ; then the minimal number of votes needed to win a district is at most  $\lceil \frac{nS_\alpha}{m} \rceil + \alpha_1$ , and at least  $\lceil \frac{nS_\alpha}{m} \rceil$ : we allocate the scores as evenly as possible among the candidates, and break the tie in favor of the winner using at most  $\alpha_1$  points. Of course, in some cases this can be improved, but it depends on  $\alpha$ , and on the divisibility of  $nS_\alpha$  and  $m$ . The following theorem uses these loose bounds to bound  $MR$  for the Borda scoring rule.

**THEOREM 3.**  $MR(V, \mathcal{D}, Borda)$  is in  $\Theta(m^2)$ .

**PROOF.** According to Lemma 2,  $L_n(Borda) = (m-1)(\lfloor \frac{n}{2} \rfloor - 1) + (m-2)(\lfloor \frac{n}{2} \rfloor + 1)$ , and  $M_n(Borda) = n(m-1)$ . We again write  $x_w$  and  $x_p$  to be the number of votes that are won by  $w$  and  $p$  (respectively) in a district where  $w$  wins. Thus,  $MR$  is

$$\frac{x_p + n(m-1) - 2(m-2)(n-1)}{x_w} + \frac{(z+1)(m-1)(m-2)n}{x_w d(w)}.$$

We can also show the score  $p$  receives in a district won by  $w$  is at least  $x_w - (m-1)$ , and at most  $x_w - 1$ . Furthermore, we observe that  $x_w$  is at least  $\lceil \frac{(m-1)(m-2)n}{m} \rceil$ , which is at least  $\frac{(m-2)^2 n}{2m}$ . Furthermore,  $x_w$  is upper bounded by  $\lceil \frac{(m-1)(m-2)n}{m} \rceil + (m-1)$ , which is at most  $\frac{m(n+2)}{2}$ . Thus,  $MR$  is upper bounded by

$$\frac{2m((m-2)n(m+1)(m-1) + (m-1))}{(m-2)^2 n}.$$

We note that the final expression has  $n$  in both the numerator and denominator and is thus  $\mathcal{O}(m^2)$ . We may similarly lower bound  $MR(V, \mathcal{D}, Borda)$  by a similar value.  $\square$

## 5.2 Copeland

When using the Copeland voting rule, one can get an undefined value for  $MR$ , as a score of 0 is possible for the winner.

*Example 1.* Let us have two identical districts, each containing 21 voters with the preference  $w \succ p \succ a$  and 20

---

### Algorithm 1 Monte-Carlo $MR$ Approximation

---

```

1: procedure EXPECTED- $MR$  ( $M, B, \varepsilon, \delta$ )
2:    $s = \lceil \frac{B^2 \cdot \ln \frac{2}{\delta}}{2\varepsilon^2} \rceil$ 
3:    $T = 0$ 
4:   for  $i = 1$  to  $s$  do
5:     Sample an election outcome  $E$  from  $M$ 
6:      $w = \arg \max_{c \in C} \text{score}(c, V)$  // The winner
7:      $s_m = \max_{c \in C} \text{score}(c, V)$  // Maximal score
8:      $R_i^{\text{samp}} = \frac{s_m}{s_w}$  // sampled  $MR$ 
9:      $T = T + R_i^{\text{samp}}$ 
10:  return  $\hat{r} = \frac{T}{s}$  // average of sampled  $MR$ s
```

---

voters who have  $p \succ w \succ a$ . A third district contains 41 voters with the preference  $p \succ w \succ a$ .  $w$  wins the 2 first districts, becoming the ultimate winner. But, looking globally,  $p$ 's Copeland score is 2, while  $w$ 's Copeland score is 0, making  $MR(V, \mathcal{D}, Copeland)$  undefined.

The Copeland score can be additively adjusted by adding to each candidate's score a fixed amount that is larger than  $m$ . However, Copeland's performance remains bad, as is captured by Theorem 4.

**THEOREM 4.** Under Copeland, the winner  $w$  may have the worst possible Copeland score, while another candidate has the best possible Copeland score.

**PROOF.** The worst possible Copeland score is  $-(m-1)$ , while the best is  $m-1$ . Taking  $z = \ell m + r$ ,  $z > 4$  and  $m > 2$ , we take  $\ell + 2$  districts, in each of them all voters rank  $w \succ p \succ \dots$ . We now take  $\ell + 1$  districts, in each of them all voters rank  $p \succ \dots \succ w$ . Finally, the rest of the districts are divided between the  $m-2$  other candidates, each candidate  $c \in C \setminus \{p, w\}$  getting at most  $\ell + 1$  districts, in which every voter ranks  $c \succ p \succ \dots \succ w$ . Since  $\ell + 2 < \frac{n}{2}$ ,  $w$  has lost to all candidates for most voters, and therefore  $w$ 's Copeland score is  $-(m-1)$ .  $p$ , on the other hand, is preferred by most voters over any other candidate, leading its score to be  $m-1$ .  $\square$

## 6. THE MISREPRESENTATION RATIO UNDER UNCERTAIN VOTES

In Section 5 we established bounds on the misrepresentation ratio by constructing pathological examples: settings where districting effects were so pronounced as to cause an extremely unpopular candidate to win the elections, despite the existence of a clearly better alternative. We now take the average-case, rather than the worst-case, approach, and ask how common are instances where misrepresentation is high.

We do this in the form of a probabilistic generative model, utilizing partial information to inform our assumptions on the general population. Any instantiation of the model is a voting domain for which we can compute the misrepresentation ratio. Thus,  $MR$  is a random variable and we evaluate the *expected*  $MR$ . A naive solution is to exhaustively search over the space of possible election outcomes; for each such outcome we can compute its probability of occurring under the generative model, and the  $MR$  value for that outcome; we can then sum the product of the two across all outcomes to get the expected  $MR$ . However, such an exhaustive search

is intractable, as the space of outcomes can be prohibitively large, especially when there are many candidates, voters and districts. We propose an alternative approximate solution, based on a Monte-Carlo algorithm. Our algorithm requires a bound on the  $MR$  for the examined domain, and such can be found for many cases (see previous section).

We assume the generative model is given in the form of a black-box, which outputs a sampled election outcome, consisting of the votes of every voter in every district. We further assume that the winner of the election can be computed in polynomial time<sup>5</sup>. Denote the generative model as  $M$ , and by  $r = E_M(MR)$  the expectation of  $MR$  under the model  $M$ . Our proposed algorithm is “probably approximately correct”: given two parameters,  $\varepsilon$  and  $\delta$ , the algorithm returns an approximation  $\hat{r}$  to  $r$ , such that with high probability  $1 - \delta$  the returned value  $\hat{r}$  is very close to  $r$ , so that  $|\hat{r} - r| \leq \varepsilon$ . The running time of the algorithm depends on  $\varepsilon$  and  $\delta$ ; it is quadratic in  $\frac{1}{\varepsilon}$  and logarithmic in  $\frac{1}{\delta}$ .

Our proposed algorithm is a Monte-Carlo algorithm, but it is only appropriate to voting rules where there is a known bound on the possible  $MR$  values.<sup>6</sup> The minimal  $MR$  value is 1 (as this is the ratio between the maximal score of any candidate and the score of a specific candidate, namely the winner). Given a bound  $H$  on the maximal  $MR$  in a domain, we refer to the  $MR$  value range as  $B = H - 1$ . The runtime of our algorithm is quadratic in  $B$ . The method is given in Algorithm 1, and we provide a proof for its correctness.

**THEOREM 5.** *The value returned by Expected- $MR$  is an  $\varepsilon, \delta$  approximation for the expected  $MR$  under  $M$ : with probability at least  $1 - \delta$  the returned value  $\hat{r}$  is within a distance  $\varepsilon$  of  $r = E_M(MR)$ , i.e.:  $|\hat{r} - r| \leq \varepsilon$ .*

**PROOF.** We note that the  $R_i^{samp}$  computed inside the loop is the  $MR$  in a specific instantiation of an election outcome  $E$  sampled from the generative model  $M$  (see Definition 2), so each  $R_i^{samp}$  is a random variable, whose expectation is  $r = E_M(MR)$  (i.e.  $E[R_i^{samp}] = r$ ). Our algorithm computes  $T = \sum_{i=1}^s R_i^{samp}$ , the sum of  $s$  i.i.d draws, each of which has a value of  $r$  in expectation, so  $E[\hat{r}] = E[\frac{T}{s}] = r$ . We use Hoeffding’s inequality [24] to show that the number of samples  $s$  that we use achieves the desired accuracy  $\varepsilon$  and confidence  $\delta$ . Hoeffding’s inequality states that if  $R_1, \dots, R_n$  are independent random variables, where each  $R_i$  is bounded so that  $R_i \in [a_i, b_i]$ , and if  $T = \sum_{i=1}^s R_i$ , then  $\Pr(|T - E[T]| \geq s\varepsilon) \leq 2 \exp\left(-\frac{2s^2\varepsilon^2}{\sum_{i=1}^n (b_i - a_i)^2}\right)$ .

$R_i^{samp}$  is the  $MR$  in an election outcome obtained under the generative model  $M$ , so  $R_i^{samp}$  is bounded in the range  $[1, H]$  (i.e., by our assumption, the  $MR$  value range is  $B = H - 1$ ). Applying Hoeffding’s bound and substituting  $\frac{B^2 \cdot \ln \frac{2}{\delta}}{2\varepsilon^2}$  for  $s$ , we get  $\Pr\left(\left|\frac{T}{s} - \frac{E[T]}{r}\right| \geq \varepsilon\right) \leq \delta$  as required.  $\square$

## 7. SIMULATIONS

We now use our Algorithm Expected- $MR$  to analyze the  $MR$  in several voting domains. We begin with a noisy version of the example domain described in the introduction.

<sup>5</sup>Not all voting rules admit a polynomial winner determination algorithm. As our algorithm samples election outcomes, its runtime in this case would not be polynomial.

<sup>6</sup>Our method is akin to Monte-Carlo methods used for analyzing voting under various forms of uncertainty [14, 2, 4, 5].




Figure 1: The effects of the noise on the  $MR$

We fix  $C = \{w, p\}$ , and the number of districts to 11: 6 districts of type “A” and 5 of type “B”, modeling heterogeneous and homogenous districts respectively. In type A districts, every voter  $v$  votes randomly with  $\Pr[v \text{ votes for } w] = \frac{1}{2} + \varepsilon$ , and  $\Pr[v \text{ votes for } p] = \frac{1}{2} - \varepsilon$ ; in type B districts,  $\Pr[v \text{ votes for } w] = \varepsilon$ ,  $\Pr[v \text{ votes for } p] = 1 - \varepsilon$ , for  $\varepsilon \in (0, \frac{1}{2})$ . Figure 1 shows the averaged  $MR$  as a function of the noise  $\varepsilon$  ( $x$ -axis). Each point is average  $MR$  obtained for many elections sampled using this probabilistic model;  $MR$  first increases as the noise  $\varepsilon$  grows, until it reaches a sweet-spot from which it drops. This indicates that for some models the model noise may not have a monotone effect on the  $MR$ .


In our second experiment, we fix the number of districts to 15, and range the number of voters in every district from 100 to 5000. We examined  $MR$  of elections with  $m \in \{3, 4, \dots, 7\}$  candidates. Figure 2a shows the averaged  $MR$  where the preference of every voter in every district is uniformly drawn from the set of all  $m!$  possible orders of candidates, under Borda scoring. The figure shows that increasing the number of voters tends to lower  $MR$ . This is not surprising, as all candidates will likely have nearly the same score. Next, we consider the Mallows model [30] for generating voter preferences, where we assume that there is a “ground truth” for every district,  $\sigma^*$  (representing the common ranking of candidates in that district) and dispersion parameter  $\phi \in (\frac{1}{2}, 1]$ . Under the Mallows model every voter compares every pair of candidates independently and ranks them correctly (according to  $\sigma^*$ ) with probability  $\phi$ . For every district,  $\sigma^*$  was drawn uniformly at random and  $\phi \sim \mathcal{U}(\frac{1}{2}, 1)$ . We used the plurality scoring rule (Figure 2b) and Copeland voting rule (Figure 2c). As predicted by our theoretical results,  $MR$  grows when there are more candidates. Under Copeland, a fixed amount of  $m$  was added each candidate’s score so that  $MR$  would be positive.

Our simulation results indicate that voting misrepresentation may occur in several natural domains. Our second experiment is “fair” in the sense that there is no preferred candidate, and yet the  $MR$  values are quite high. Also, our theoretical results agree with experiments in some natural domains.

## 8. CONCLUSIONS

This work analyzes district-based elections. We demonstrate the representability issues that arise in such elections, and show tight bounds on misrepresentation. We further show that misrepresentation is a common occurrence under various natural voter distributions, and that its effect may





(a) *MR* where the preferences are uniformly drawn, under Borda scoring rule (b) *MR* where the preferences are drawn by a Mallows model, under plurality (c) *MR* where the preferences are drawn by a Mallows model, under Copeland voting rule

Figure 2: Simulations results

not diminish even when the number of voters is large.

District based elections tend to under-represent smaller parties; this is a long observed phenomenon (and, in some countries, a welcome stabilizing feature). However, we do not focus on smaller parties, but rather show that the preferences of large majorities may be completely unrepresented (the UK elections of 1951, where the Labour party with 48.8% support lost the elections, and the Conservative party with 44.3% not only beat it, but had a strong parliamentary majority, is just one example of these occurrences).

Research into the institutional bias in voting procedures, beyond control issues, is one which we think deserves more attention by the computational social choice community. Very few election systems in the world are proportional, and the effect this has on the expression of voters' views has mostly focused (in political science research) on how small minorities are hurt. As our analysis shows, large majorities may also be affected. Further research is needed with regards to other voting methods. Moreover, further and complementary concepts may be developed, indicating unfairness, lack of representation and other problems with various voting procedures (parliamentary entrance bounds, common in some countries, are an obvious candidate for such directions). In addition, while we have focused on an outcome of a single winner, a coalitional analysis of district settings may also be of interest.

## Acknowledgements

This work was supported in part by NSERC grant 482671.

## REFERENCES

- [1] H. Aziz, Y. Bachrach, E. Elkind, and M. Paterson. False-name manipulations in weighted voting games. *Journal of Artificial Intelligence Research*, pages 57–93, 2011.
- [2] Y. Bachrach, N. Betzler, and P. Faliszewski. Probabilistic possible winner determination. In *AAAI*, volume 10, pages 697–702, 2010.
- [3] Y. Bachrach, E. Elkind, and P. Faliszewski. Coalitional voting manipulation: A game-theoretic perspective. In *IJCAI Proceedings-International Joint Conference on Artificial Intelligence*, volume 22, page 49, 2011.
- [4] Y. Bachrach, R. Meir, M. Feldman, and M. Tennenholtz. Solving cooperative reliability games. *arXiv preprint arXiv:1202.3700*, 2012.
- [5] Y. Bachrach and N. Shah. Reliability weighted voting games. In *Algorithmic Game Theory*, pages 38–49. Springer, 2013.
- [6] J. J. Bartholdi, C. A. Tovey, and M. A. Trick. The computational difficulty of manipulating an election. *SCW*, 6(3):227–241, 1989.
- [7] D. Butler. The electoral process the redrawing of parliamentary boundaries in Britain. *British Elections and Parties Yearbook*, 2(1), 1992.
- [8] E. J. Engstrom. Stacking the states, stacking the house: The partisan consequences of congressional redistricting in 19th century. *APSR*, 100:419–427, 2006.
- [9] G. Erdélyi, E. Hemaspaandra, and L. A. Hemaspaandra. More natural models of electoral control by partition. In *ADT*, pages 396–413, 2015.
- [10] R. S. Erikson. Malapportionment, gerrymandering, & party fortunes in congressional elections. *APSR*, 66(4):1234–1245, 1972.
- [11] P. Faliszewski, E. Hemaspaandra, and L. A. Hemaspaandra. The complexity of bribery in elections. In *AAAI*, volume 6, pages 641–646, 2006.
- [12] P. Faliszewski, E. Hemaspaandra, and L. A. Hemaspaandra. Using complexity to protect elections. *CACM*, 53(11):74–82, 2010.
- [13] P. Faliszewski, E. Hemaspaandra, L. A. Hemaspaandra, and J. Rothe. Llull and copeland voting computationally resist bribery and constructive control. *Journal of Artificial Intelligence Research*, pages 275–341, 2009.
- [14] S. S. Fatima, M. Wooldridge, and N. R. Jennings. A linear approximation method for the shapley value. *Artificial Intelligence*, 172(14):1673–1699, 2008.
- [15] M. Feix, D. Lepelley, V. Merlin, J. Rouet, and L. Vidu. Majority efficient representation of the citizens in a federal union. Technical report, Université de la Réunion, de Caen, d’Orleans, 2008.
- [16] M. R. Feix, D. Lepelley, V. R. Merlin, and J. Rouet. The probability of conflicts in a U.S. presidential type election. *Economic Theory*, 23(2):227–257, 2004.
- [17] D. S. Felsenthal and M. Machover. Minimizing the mean majority deficit: The 2nd square-root rule. *Math. Soc. Sci.*, 37(1):25–37, 1999.
- [18] J. N. Friedman and R. T. Holden. Towards a theory of optimal partisan gerrymandering. Technical report,



- Harvard University, 2005.
- [19] M. Gallagher. Proportionality, disproportionality and electoral systems. *Electoral Studies*, 10(1):33–51, 1991.
- [20] A. Gelman, J. N. Katz, and F. Tuerlinckx. The mathematics and statistics of voting power. *Statistical Science*, 17(4):420–435, 2002.
- [21] M. Goemans and T. Rothvoß. Polynomiality for bin packing with a constant number of item types. In *SODA*, pages 830–839, 2014.
- [22] B. Grofman, W. Koetzle, and T. Brunell. An integrated perspective on the three potential sources of partisan bias: Malapportionment, turnout differences, & the geographic distribution of party vote shares. *Electoral Studies*, 16(4):457–470, 1997.
- [23] E. Hemaspaandra, L. A. Hemaspaandra, and J. Rothe. Anyone but him: The complexity of precluding an alternative. *AIJ*, 171(5–6):255–285, 2007.
- [24] W. Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association*, 58(301):13–30, 1963.
- [25] S. Issacharoff. Gerrymandering & political cartels. *Harvard Law Review*, 116(2):593–648, 2002.
- [26] K. Jansen, S. Kratsch, D. Marx, and I. Schlotter. Bin packing with fixed number of bins revisited. *JCSC*, 79(1):39–49, 2013.
- [27] R. Lahrach and V. Merlin. *Electoral Systems: Paradoxes, Assumptions, and Procedures*, chapter Which Voting Rule Minimizes the Probability of the Referendum Paradox? Lessons from French Data, pages 129–150. Springer, 2011.
- [28] R. Lahrach and V. R. Merlin. Assessing the probability of the referendum paradox: The french local election case. Voting Power in Practice workshop, 2010.
- [29] D. Lublin. *The Paradox of Representation: Racial Gerrymandering and Minority Interests in Congress*. Princeton University Press, 1997.
- [30] C. Mallows. Non-null ranking models. I. *Biometrika*, pages 114–130, 1957.
- [31] N. Miller. *Electoral Systems: Paradoxes, Assumptions & Procedures*, chapter Election Inversions by the U.S. Electoral College, pages 93–127. Springer, 2011.
- [32] N. Miller. The house size effect and the referendum paradox in u.s. presidential elections. *Electoral Studies*, 35:265–271, September 2014.
- [33] H. Nurmi. *Voting Paradoxes and How to Deal with Them*. Springer-Verlag, 1999.
- [34] G. Owen and B. Grofman. Optimal partisan gerrymandering. *Political Geography Quarterly*, 7(1):5–22, January 1988.
- [35] A. D. Procaccia, J. S. Rosenschein, and A. Zohar. Multi-winner elections: Complexity of manipulation, control & winner-determination. In *IJCAI*, pages 1476–1481, 2007.
- [36] P. Schuck. The thickest thicket: Partisan gerrymandering and judicial regulation of politics. *Columbia Law Review*, 87(7):1325–1384, 1987.
- [37] W. Stille. *Solution Techniques for specific Bin Packing Problems with Applications to Assembly Line Optimization*. PhD thesis, TU Darmstadt, June 2008.
- [38] A. Tangian. Computational application of the mathematical theory of democracy to Arrow’s impossibility theorem (how dictatorial are Arrow’s dictators?). *SCW*, 35(1):129–161, 2010.
- [39] L. Xia. Computing the margin of victory for various voting rules. In *EC*, pages 982–999, 2012.
- [40] L. Xia, M. Zuckerman, A. D. Procaccia, V. Conitzer, and J. S. Rosenschein. Complexity of unweighted coalitional manipulation under some common voting rules. In *IJCAI*, pages 348–353, 2009.
- [41] M. Zuckerman, A. D. Procaccia, and J. S. Rosenschein. Algorithms for the coalitional manipulation problem. In *SODA*, pages 277–286, 2008.